

## Improved time-variable gravity modelling using monthly COST-G models for precise orbit determination of low Earth orbiting satellites

Adrian Jäggi<sup>1</sup>, Heike Peter<sup>2</sup>, Ulrich Meyer<sup>1</sup>

<sup>1</sup>University of Bern, Astronomical Institute, Switzerland <sup>2</sup>Positim UG, Germany

2022 Ocean Surface Topography Science Team Meeting October 31 – November 4, 2022, Venice, Italy






### Combination Service for Time-variable Gravity Fields (COST-G)




#### Introduction to COST-G



Improved and consolidated product integrating the strengths of all ACs




#### **Workflow of COST-G**






#### **GRACE-FO** operational combined monthly gravity fields



Flawless and uninterrupted operational combination with a latency < 2-3 months.



#### Weighted combination and validation of the Combined Solution





### https://cost-g.org/



For background information on COST-G and links to products take a look at: https://cost-g.org/

and visit the poster POD2022-004

#### Welcome to COST-G

The International Combination Service for Time-variable Gravity Fields (COST-G) is a product center of the International Gravity Field Service (IGFS) and is dedicated to the combination of monthly global gravity field models. COST-G stems from the activities of the former H2020 project European Gravity Service for Improved Emergency Management (EGSIEM) and is further developed within the follow-up project Global Gravity-Based Groundwater Product (G3P), which is funded from the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement no. 870353 (funding period 2020-2022).

Please use the top menu to visit the various parts of our website!

Best regards, Your COST-G Team.

#### **Latest News**

April 14th 2022

We have a new publication online:
COST-G gravity field models for precise orbit determination of Low Earth Orbiting Satellites.

December 17th 2021

Precise orbit determination (POD) of Low Earth Orbiters (LEOs) depends on the precise knowledge of the Earth's gravity field Peter H, Meyer U, Lasser M, Jäggi A (2022): COST-G gravity field models for precise orbit determination of Low Earth Orbiting Satellites.
Advances in Space Research (69), 12, 4155-4168.

doi: 10.1016/j.asr.2022.04.005





#### **2022 OSTST Meeting**

# Fitted Signal Model (FSM) for operational LEO-POD



#### RMS of differences (over land, 300 km Gauss): FSM - monthly gravity fields





Operational precise orbit determination (POD) of low Earth orbiters (LEO) relies on a Earth gravity model including time-variable gravity (TVG).


The EIGEN-GRGS-RL04 model (green) has been the standard for LEO-POD of altimeter satellites, but the extrapolation to the GRACE-FO period reveals large prediction errors.

For comparison, a model fitted to COST-G GRACE-FO gravity fields is shown (red).





#### Polar mass trend (no filter)




Surprisingly, the reason for the prediction error in the EIGEN-GRGS-RL04 model (green) seems not to be in regions with strong mass trends.





#### Hydrological cycle in large river basins (300 km Gauss)



The time-series of monthly GRACE gravity field solutions was fitted in yearly batches for the EIGEN-GRGS-RL04 model.

While the fit in the GRACE period is very good, the extrapolation of the last of these batches leads to large errors in river basins with strong non-seasonal variations.






## Application to Sentinel orbit POD



#### Sentinel - 3B (altitude 811 km) orbit determination





The carrier phase RMS of dynamic Sentinel-3B satellite orbits (orbit altitude 811 km) based on monthly GRACE-FO gravity fields (green) or different fitted signal models reveals the benefit of up-to-date models. All models were truncated at max. degree/order 90.





#### Impact of fit period on LEO-POD (Sentinel-3B, altitude 811 km)





Carrier phase residuals of Sentinel-3B orbits (811 km orbit altitude) confirm the sensitivity on the data period that entered the model.





## Independent orbit validation



#### **SLR-validation Sentinel-3B**

#### Data: Year 2020, Sentinel-3B, SLR validation, 12 stations (cm)

| Gravity field model | Mean (cm) | RMS (cm) | Standard deviation (cm) |
|---------------------|-----------|----------|-------------------------|
| DEIGEN120           | 0.29      | 1.01     | 0.97                    |
| DEIGEN90            | 0.29      | 1.01     | 0.97                    |
| D90MONTHLY          | 0.28      | 0.91     | 0.87                    |
| D90MODEL2012        | 0.28      | 0.92     | 0.88                    |
| RDEIGEN120          | 0.31      | 0.91     | 0.85                    |
| RDEIGEN90           | 0.31      | 0.91     | 0.85                    |
| RD90MONTHLY         | 0.31      | 0.88     | 0.82                    |

The limited max. degree does not negatively affect LEO POD (S3B)

LEO POD profits from monthly gravity fields

The fitted signal models perform close to the monthly gravity fields

Reduced dynamic LEO POD is less sensitive to model deficiencies.






### Outlook



#### **Extension of COST-G FSM for REPRO purposes**




Extension of the COST-G FSM to cover the whole GRACE/GRACE-FO period:

- Fit of GRACE monthly models in yearly batches
- Continuity conditions between individual batches
- Fit of GRACE-FO monthly models in one batch to allow for prediction.



#### **Consistency with monthly models**




The monthly GRACE gravity fields were screened prior to the fit of the extended COST-G FSM => outliers indicate monthly solutions of inferior quality.



# Where to get the COST-G fitted signal models?



### http://icgem.gfz-potsdam.de/series



User-defined points

3D Visualisation

Static Models

**Temporal Models** 

Trend & Amplitude

**Spherical Harmonics** 

Evaluation

Spectral domain

**GNSS Leveling** 



#### **Gravity Field Solutions for dedicated Time Periods**

The following gravity field time series are presently available:

|   |             |  | nce Data System centers CSR, GFZ and JPL collapse  |  |  |
|---|-------------|--|--|--|--|
|   |             |  | Center for Space Research at University of Texas, Austin   |  |  |
| CSR Release 05 monthly  |             |  | UTCSR Level-2 Processing Standards Document, Rev 4.0 May 29, 2012                                |  |  |
| CSR Release 06 DOI monthly  |             |  | UTCSR Level-2 Processing Standards Document, Rev 5.0 April 18, 2018                              |  |  |
| SR Release 06 (GFO) DOI monthly UTCSR Level-2 Processing Standards Document, V 1.1. |             |  | UTCSR Level-2 Processing Standards Document, V 1.1 June 6, 2019                                  |  |  |
| - GFZ   |             |  | Helmholtz Centre Potsdam German Research Centre for Geosciences                                  |  |  |
|   | monthly     | weekly   | GFZ GRACE Level-2 Processing, Revised Edition, January 2013                                      |  |  |
| GFZ Release 06 DOI monthly  |             |  | GFZ GRACE Level-2 Processing Standards Document for Level-2 Products, Rev. 1.0, October 26, 2018 |  |  |
| DOI   | monthly     |  | GFZ GRACE Level-2 Processing Standards Document for Level-2 Products, Rev. 1.0, June 3, 2019     |  |  |
| - JPL JPL Release 05 monthly  |             |  | Jet Propulsion Laboratory  |  |  |
|   |             |  | JPL Level-2 Processing Standards Document, Release 05.1 November 3, 2014                         |  |  |
| DOI   | monthly     |  | JPL Level-2 Processing Standards Document, Release 06.0 June 1, 2018                             |  |  |
| DOI   | monthly     |  | JPL Level-2 Processing Standards Document, v 1.0 May 28, 2019                                    |  |  |
|   | DOI DOI DOI | DOI monthly DOI monthly  monthly DOI monthly DOI monthly monthly DOI monthly | DOI monthly  monthly weekly  DOI monthly  monthly  monthly  monthly  monthly  DOI monthly        |  |  |

The processing standards to generate the GRACE Level-2 products of CSR, GFZ and JPL are also available in the Document Section of the GRACE archives at GFZ ISDC or JPL PO.DAAC

| C | OST-G (Internati | onal Comb | ination Servic | e for Time-variable Gravity F |
|---|------------------|-----------|----------------|-------------------------------|
|   | DSM              |           | quarterly      | Deterministic Signal Model    |
|   | Grace            | וטע       | monthly        |                               |
|   | Grace-FO         | DOI       | monthly        |                               |
|   | Swarm            | DOI       | monthly        |                               |

icgem (at) gfz-potsdam.de

The COST-G fitted signal model is available in the ICGEM.2-format from the International Center for Global Earth Models (ICGEM).

The COST-G FSM is updated quarterly with the newest combined monthly GRACE-FO gravity fields.



#### **Summary**

- COST-G GRACE-FO combined Level-2 products are made available with a latency of approx. 3 months at ICGEM.
- A revised weighting scheme has been found that is in good accordance with the noise assessment of the individual AC solutions.
- The combined solution is shown to outperform individual AC solutions in terms of the noise assessment over the oceans.
- Fitted Signal Models (FSM) are provided by COST-G to support operational LEO-POD activities.
- FSM are fitted to GRACE-FO monthly combined solutions and updated quarterly
- Outlook: extension of FSM to GRACE period is currently on-going



### Thanks a lot for your attention

